

## DAFTAR PUSTAKA

- Agarwal, Pathak D., Bhutani, Kapoor, & Kant. (2017). Review on *Camellia sinensis* –Nature’s Gift. *International Journal of Pharmacognosy and Phytochemical Research*, 9(08). <https://doi.org/10.25258/phyto.v9i08.9619>
- Agustina, E., Andiarna, F., Lusiana, N., Purnamasari, R., & Irfan Hadi, M. (2018). Identifikasi Senyawa Aktif dari Ekstrak Daun Jambu Air (*Syzygium aqueum*) dengan Perbandingan Beberapa Pelarut pada Metode Maserasi. *BIOTROPIC The Journal of Tropical Biology*, 2(2). <http://jurnalsaintek.uinsby.ac.id/index.php/biotropic>
- Aly, A. A., Sayed, S. M., Abdelhafez, E. S. M. N., Abdelhafez, S. M. N., Abdelzaher, W. Y., Raslan, M. A., Ahmed, A. E., Thabet, K., El-Reedy, A. A. M., Brown, A. B., & Bräse, S. (2020). New quinoline-2-one/pyrazole derivatives; design, synthesis, molecular docking, anti-apoptotic evaluation, and caspase-3 inhibition assay. *Bioorganic Chemistry*, 94. <https://doi.org/10.1016/J.BIOORG.2019.103348>
- Arjunan, S., Gan, S. U., Choolani, M., Raj, V., Lim, J., Biswas, A., Bongso, A., & Fong, C. Y. (2020). Inhibition of growth of Asian keloid cells with human umbilical cord Wharton’s jelly stem cell-conditioned medium. *Stem Cell Research and Therapy*, 11(1). <https://doi.org/10.1186/s13287-020-01609-7>
- Berman, B., Maderal, A., & Raphael, B. (2017). Keloids and hypertrophic scars: Pathophysiology, classification, and treatment. In *Dermatologic Surgery* (Vol. 43, pp. S3–S18). Lippincott Williams and Wilkins. <https://doi.org/10.1097/DSS.0000000000000819>
- Binic, I. (2015). Keloids and hypertrophic scars. In *European Handbook of Dermatological Treatments, Third Edition* (pp. 455–464). Springer Berlin Heidelberg. [https://doi.org/10.1007/978-3-662-45139-7\\_47](https://doi.org/10.1007/978-3-662-45139-7_47)
- Chen, L., Zhang, G., He, J., Guan, J., Pan, C., Mi, W., & Wang, Q. (2010). New naphthoquinone from the root of *Lygodium japonicum* (Thunb.) Sw. *Journal of Natural Medicines*, 64(1), 114–116. <https://doi.org/10.1007/s11418-009-0376-y>
- Chen, Z., Zhang, N., Chu, H. Y., Yu, Y., Zhang, Z. K., Zhang, G., & Zhang, B. T. (2020). Connective Tissue Growth Factor: From Molecular Understandings to Drug Discovery. In *Frontiers in Cell and Developmental Biology* (Vol. 8). Frontiers Media S.A. <https://doi.org/10.3389/fcell.2020.593269>

- Cummings, B. S., & Schnellmann, R. G. (2002). *Cisplatin-Induced Renal Cell Apoptosis: Caspase 3-Dependent and-Independent Pathways*. <http://jpet.aspetjournals.org>
- El Naga, R. N. A., Azab, S. S., El-Demerdash, E., Shaarawy, S., El-Merzabani, M., & Ammar, E. S. M. (2013). Sensitization of TRAIL-induced apoptosis in human hepatocellular carcinoma HepG2 cells by phytochemicals. *Life Sciences*, 92(10), 555–561. <https://doi.org/10.1016/j.lfs.2013.01.017>
- Ellis, S., Lin, E. J., & Tartar, D. (2018). Immunology of Wound Healing. In *Current Dermatology Reports* (Vol. 7, Issue 4, pp. 350–358). Current Medicine Group LLC 1. <https://doi.org/10.1007/s13671-018-0234-9>
- Gallucci, R. M., Lee, E. G., & Tomasek, J. J. (2006). IL-6 modulates alpha-smooth muscle actin expression in dermal fibroblasts from IL-6-deficient mice. *Journal of Investigative Dermatology*, 126(3), 561–568. <https://doi.org/10.1038/sj.jid.5700109>
- Ghazizadeh, M., Tosa, M., Shimizu, H., Hyakusoku, H., & Kawanami, O. (2007). Functional implications of the IL-6 signaling pathway in keloid pathogenesis. *Journal of Investigative Dermatology*, 127(1), 98–105. <https://doi.org/10.1038/sj.jid.5700564>
- Haghighatdoost, F., & Hariri, M. (2019). The effect of green tea on inflammatory mediators: A systematic review and meta-analysis of randomized clinical trials. In *Phytotherapy Research* (Vol. 33, Issue 9, pp. 2274–2287). John Wiley and Sons Ltd. <https://doi.org/10.1002/ptr.6432>
- Haly, Q., Sartika, C. R., Haifa, R., Karina, N., Rahmawati, V., & Naura, N. F. (2022). Combination treatment of Umbilical Cord-derived Mesenchymal Stem Cell (UC-MSC) and Umbilical Cord Mesenchymal Stem Cell-derived Conditioned Medium (UCMSC-CM) for Keloid Post Chronic Burn Injury: A Case Report. In *Malaysian Journal of Medicine and Health Sciences* (Vol. 18, Issue 1).
- Johnson, B. Z., Stevenson, A. W., Prêle, C. M., Fear, M. W., & Wood, F. M. (2020). The role of IL-6 in skin fibrosis and cutaneous wound healing. *Biomedicine*, 8(5). <https://doi.org/10.3390/BIOMEDICINES8050101>
- Kari, S., Subramanian, K., Altomonte, I. A., Murugesan, A., Yli-Harja, O., & Kandhavelu, M. (2022). Programmed cell death detection methods: a systematic review and a categorical comparison. *Apoptosis*, 27(7–8), 482. <https://doi.org/10.1007/S10495-022-01735-Y>

- Karppinen, S. M., Heljasvaara, R., Gullberg, D., Tasanen, K., & Pihlajaniemi, T. (2019). Toward understanding scarless skin wound healing and pathological scarring [version 1; peer review: 2 approved]. In *F1000Research* (Vol. 8). F1000 Research Ltd. <https://doi.org/10.12688/f1000research.18293.1>
- Kendall, R. T., & Feghali-Bostwick, C. A. (2014). Fibroblasts in fibrosis: Novel roles and mediators. *Frontiers in Pharmacology*, 5 MAY. <https://doi.org/10.3389/fphar.2014.00123>
- Kendig, E. L., Le, H. H., & Belcher, S. M. (2010). Defining hormesis: Evaluation of a complex concentration response phenomenon. In *International Journal of Toxicology* (Vol. 29, Issue 3, pp. 235–246). <https://doi.org/10.1177/1091581810363012>
- Khan, N., Afaq, F., Saleem, M., Ahmad, N., & Mukhtar, H. (2006). Targeting multiple signaling pathways by green tea polyphenol (-)-epigallocatechin-3-gallate. In *Cancer Research* (Vol. 66, Issue 5, pp. 2500–2505). <https://doi.org/10.1158/0008-5472.CAN-05-3636>
- Koch, W., Zagórska, J., Marzec, Z., & Kukula-Koch, W. (2019). Applications of tea (*Camellia sinensis*) and its active constituents in cosmetics. In *Molecules* (Vol. 24, Issue 23). MDPI AG. <https://doi.org/10.3390/molecules24234277>
- Li, M., Xu, J., Shi, T., Yu, H., Bi, J., & Chen, G. (2016). Epigallocatechin-3-gallate augments therapeutic effects of mesenchymal stem cells in skin wound healing. *Clinical and Experimental Pharmacology and Physiology*, 43(11), 1115–1124. <https://doi.org/10.1111/1440-1681.12652>
- Luo, S. M. D., Benathan, M. Ph. D., Raffoul, W. M. D., Panizzon, R. G. M. D., & Egloff, D. V. M. D. (2001). Abnormal Balance Between Proliferation and Apoptosis Cell Death in Fibroblasts Derived from Keloid Lesions. *Plastic and Reconstructive Surgery*, 107(1), 87–95.
- Luo, X., Lu, F., Yin, Z., Zhou, Z., Wang, Z., & Zhang, H. (2023). Hormetic effects of EGC and EGCG on CES1 activity and its rescue from oxidative stress in rat liver S9. *Chemico-Biological Interactions*, 382, 110612. <https://doi.org/10.1016/J.CBI.2023.110612>
- Mahboub, F. A., & Khorshid, F. A. (2010). The Role of Green Tea Extract on the Proliferation of Human Ovarian Cancer Cells (in vitro) Study. *International Journal of Cancer Research*, 6(2), 78–88. <https://doi.org/DOI: 10.3923/ijcr.2010.78.88>

- Mari, W., Alsabri, S. G., Tabal, N., Younes, S., Sherif, A., & Simman, R. (2015). Novel Insights on Understanding of Keloid Scar: Article Review. In *Journal of the American College of Clinical Wound Specialists* (Vol. 7, Issues 1–3, pp. 1–7). Elsevier Inc. <https://doi.org/10.1016/j.jccw.2016.10.001>
- Messadi, D. v., Doung, H. S., Zhang, Q., Kelly, A. P., Tuan, T. L., Reichenberger, E., & Le, A. D. (2004). Activation of NFκB signal pathways in keloid fibroblasts. *Archives of Dermatological Research*, 296(3), 125–133. <https://doi.org/10.1007/s00403-004-0487-y>
- Meutia Sari, L. (2018). APOPTOSIS: MEKANISME MOLEKULER KEMATIAN SEL (Tinjauan Pustaka) APOPTOSIS: MOLECULAR MECHANISMS OF CELLULAR DEATH (Literature Review). *Cakradonya Dental Journal*, 10(2), 65–70. <http://www.jurnal.unsyiah.ac.id/CDJ>
- Millipore. (2018). *Certificate of Analysis / Protocol Human IL-6 ELISA Kit*.
- Moodley, Y. P., Misso, N. L. A., Scaffidi, A. K., Fogel-Petrovic, M., McAnulty, R. J., Laurent, G. J., Thompson, P. J., & Knight, D. A. (2003). Inverse effects of interleukin-6 on apoptosis of fibroblasts from pulmonary fibrosis and normal lungs. *American Journal of Respiratory Cell and Molecular Biology*, 29(4), 490–498. <https://doi.org/10.1165/RCMB.2002-0262OC>
- Motoki, T. H. C., Isoldi, F. C., Brito, M. J. de A., Filho, A. G., & Ferreira, L. M. (2019). Keloid negatively affects body image. *Burns*, 45(3), 610–614. <https://doi.org/10.1016/j.burns.2018.10.009>
- Nangole, F. W., & Agak, G. W. (2019). Keloid pathophysiology: fibroblast or inflammatory disorders?: Pathophysiology of Keloids. In *JPRAS Open* (Vol. 22, pp. 44–54). Elsevier Ltd. <https://doi.org/10.1016/j.jpra.2019.09.004>
- Olaitan P.B, Olabanji J.K, Oladele A.O, & Oseni G.A. (2013). Symptomatology of Keloids in Africans. *Leone Journal of Biomedical Research*, 5(1), 29–33.
- Plikus, M. V., Wang, X., Sinha, S., Forte, E., Thompson, S. M., Herzog, E. L., Driskell, R. R., Rosenthal, N., Biernaskie, J., & Horsley, V. (2021). Fibroblasts: Origins, definitions, and functions in health and disease. In *Cell* (Vol. 184, Issue 15, pp. 3852–3872). Elsevier B.V. <https://doi.org/10.1016/j.cell.2021.06.024>
- Purnama, H., & Ratnawulan, S. (2017). *REVIEW SISTEMATIK: PROSES PENYEMBUHAN DAN PERAWATAN LUKA*.

- Sato, C., Yamamoto, Y., Funayama, E., Furukawa, H., Oyama, A., Murao, N., Hosono, H., Kawakubo, K., Sakamoto, N., & Ohnishi, S. (2018). Conditioned medium obtained from amnion-derived mesenchymal stem cell culture prevents activation of keloid fibroblasts. *Plastic and Reconstructive Surgery*, *141*(2), 390–398. <https://doi.org/10.1097/PRS.00000000000004068>
- Seifert, O., & Mrowietz, U. (2009). Keloid scarring: Bench and bedside. In *Archives of Dermatological Research* (Vol. 301, Issue 4, pp. 259–272). <https://doi.org/10.1007/s00403-009-0952-8>
- Shaheen, A. A. (2017). Risk Factors of Keloids: A Mini Review. *Austin Journal of Dermatology*, *4*(2). <https://doi.org/10.26420/austinjdermatolog.2017.1074>
- Sihite, B. A. (2013). *Ekstrak Metanol Daun Kelor Menurunkan Ekspresi BCL-2, TRAIL-R1, dan Kadar Caspase-3 Jaringan Kolon Tikus yang Diinduksi DMBA*.
- Singh, B. N., Shankar, S., & Srivastava, R. K. (2011). Green tea catechin, epigallocatechin-3-gallate (EGCG): Mechanisms, perspectives and clinical applications. In *Biochemical Pharmacology* (Vol. 82, Issue 12, pp. 1807–1821). Elsevier Inc. <https://doi.org/10.1016/j.bcp.2011.07.093>
- Sinto, L. (2018). *Scar Hipertrofik dan Keloid: Patofisiologi dan Penatalaksanaan* (Vol. 45, Issue 1).
- Sugiyono. (2013). Metode Penelitian Pendidikan Pendekatan Kuantitatif, Kualitatif, dan R&D. In *Metode Penelitian Pendidikan Pendekatan Kuantitatif, Kualitatif, dan R&D*. [https://doi.org/10.1016/S0084-3970\(08\)79262-4](https://doi.org/10.1016/S0084-3970(08)79262-4)
- Sun, L., Zhang, Y., Zhang, W., Lai, X., Li, Q., Zhang, L., & Sun, S. (2020). Green tea and black tea inhibit proliferation and migration of HepG2 cells via the PI3K/Akt and MMPs signalling pathway. *Biomedicine & Pharmacotherapy*, *125*, 109893. <https://doi.org/10.1016/J.BIOPHA.2020.109893>
- Syed, F., Bagabir, R. A., Paus, R., & Bayat, A. (2013). Ex vivo evaluation of antifibrotic compounds in skin scarring: EGCG and silencing of PAI-1 independently inhibit growth and induce keloid shrinkage. *Laboratory Investigation*, *93*(8), 946–960. <https://doi.org/10.1038/labinvest.2013.82>
- Tsai, C.-H., & Ogawa, R. (2019). Keloid research: current status and future directions. *Scars, Burns & Healing*, *5*, 205951311986865. <https://doi.org/10.1177/2059513119868659>

- Wiegand, C., Hipler, U. C., Elsner, P., & Tittelbach, J. (2021). Keratinocyte and fibroblast wound healing in vitro is repressed by non-optimal conditions but the reparative potential can be improved by water-filtered infrared A. *Biomedicines*, 9(12). <https://doi.org/10.3390/biomedicines9121802>
- Wintoko, R., & Yadika, A. D. N. (2020). Manajemen Terkini Perawatan Luka. *JK UNILA*, 4(2).
- Xu, F. W., Lv, Y. L., Zhong, Y. F., Xue, Y. N., Wang, Y., Zhang, L. Y., Hu, X., & Tan, W. Q. (2021). Beneficial effects of green tea EGCG on skin wound healing: A comprehensive review. In *Molecules* (Vol. 26, Issue 20). MDPI. <https://doi.org/10.3390/molecules26206123>
- Xu, G., & Shi, Y. (2007). Apoptosis signaling pathways and lymphocyte homeostasis. *Cell Research*, 17(9), 759–771. <https://doi.org/10.1038/cr.2007.52>
- Xu, J., Yang, E., Yu, N.-Z., Wang, Y.-B., & Long, X. (2017). The radiation therapy in keloids treatment: a comprehensive review of pathomechanism, damage mechanisms and cellular response. *Plastic and Aesthetic Research*, 4(0), null. <https://doi.org/10.20517/2347-9264.2017.24>
- Ye, Q., Wang, S. J., Chen, J. Y., Rahman, K., Xin, H. L., & Zhang, H. (2015). Medicinal plants for the treatment of hypertrophic scars. In *Evidence-based Complementary and Alternative Medicine* (Vol. 2015). Hindawi Limited. <https://doi.org/10.1155/2015/101340>
- Zhang, Q., Yamaza, T., Kelly, A. P., Shi, S., Wang, S., Brown, J., Wang, L., French, S. W., Shi, S., & Le, A. D. (2009). Tumor-like stem cells derived from human keloid are governed by the inflammatory niche driven by IL-17/IL-6 axis. *PLoS ONE*, 4(11). <https://doi.org/10.1371/journal.pone.0007798>